

South West London Inflammatory Bowel Disease Pathway

Developed and agreed by the SWL IBD Medicines Optimisation Clinical Network

Version 4.2 (09/09/2020)

Based on recommendations made by the
SWL IBD Medicines Optimisation network meeting held on 28/02/2019
and subsequent amendment ratified at SWL Medicines Optimisation Group on 26/03/2020


Contents:

Presenting with IBD symptoms.....	2
CROHN'S DISEASE - Pathway 1: Inducing and maintaining remission.....	3
CROHN'S DISEASE - Pathway 2: Immuno-modulators.....	4
CROHN'S DISEASE - Pathway 3: High Cost Drugs.....	5
Pathway 3A: Severe active Crohn's Disease.....	6
Pathway 3B: Active fistulising Crohn's Disease.....	7
ULCERATIVE COLITIS - Pathway 1: Inducing and maintaining remission.....	8
ULCERATIVE COLITIS - Pathway 2: Immuno-modulators.....	9
ULCERATIVE COLITIS - Pathway 3: High Cost Drugs.....	10
Pathway 3A: Moderately to severely active ulcerative colitis managed in outpatients.....	11
Pathway 3B: Acute exacerbation of severely active ulcerative colitis with hospitalisation.....	12
High Cost Drugs - Contraindications, cautions and information about malignancies.....	13
High Cost Drugs - Dose escalation strategy.....	14
References.....	15
Version control.....	16

This pathway provides guidance to healthcare professionals in primary and secondary care for adult patients only. However, always use clinical judgement when a patient presents with IBD symptoms.

SWL IBD Pathway – Presenting with IBD symptoms

(References: 1, 2, 3, 4 with local agreements) (09/09/2020)



IBD contact details for healthcare professionals:	Tel:	Switchboard:	E-mail:
Epsom and St. Helier Hospital (Epsom site)	IBD helpline: 01372 735363 Secretary: 01372 735129	02082962000 (ext:5129)	esth.cnsgastroegh@nhs.net ugi-egh@nhs.net
Epsom and St. Helier Hospital (St. Helier site)	IBD helpline: 07831 120969	02082962000 (ext:2340)	ugi-sth@nhs.net
Kingston Hospital	IBD helpline: 020 8934 2760	020 8546 7711	khft.ibdadviseline@nhs.net
St. George's Hospital	IBD helpline: 020 8725 2996	020 8672 1255	stgh-tr.ibdadviseline@nhs.net
Croydon University Hospital	IBD helpline: 020 8401 3000	020 8401 3000 (ext:4484)	ch-tr.ibdcuh@nhs.net

SWL IBD Pathway CROHN'S DISEASE - Pathway 1: Inducing and maintaining remission

(Reference: NICE Guideline NG129⁵) (09/09/2020)

Advice

Discuss treatment options and monitoring. Give advice on:

- Smoking cessation
- Patient experience
- Medicines adherence
- Fertility

If appropriate give information on:

- Diet and nutrition
- Fertility, pregnancy and sexual relationships
- Prognosis
- Side effects of treatment

- Cancer risk
- Surgery
- Support groups

INDUCING REMISSION

First presentation or single inflammatory exacerbation in a 12 months period

- 2 inflammatory exacerbations in a 12 months period or
- steroid dependent (glucocorticosteroid dose cannot be tapered)

Monotherapy with conventional glucocorticosteroid:

- Prednisolone or
- Methylprednisolone or
- Hydrocortisone IV

Patients who decline, cannot tolerate or have a contraindication to conventional glucocorticosteroids:

- Budesonide (disto-ileal, ileo-caecal or right-sided colonic disease) or
- Aminosalicylate (*unlicensed*) - less effective than budesonide

Explain that budesonide and mesalazine are less effective than conventional glucocorticosteroids but may have fewer side-effects.

Do not offer:

- Budesonide or mesalazine for severe presentations or exacerbations
- Azathioprine, 6-mercaptopurine or methotrexate as monotherapy to induce remission

Surgery (distal ileum):

Consider if disease is limited to distal ileum early in the course of the disease as an alternative to medical treatment

Balloon dilatation:

Consider if strictures

Consider immuno-modulators

CROHN'S DISEASE – Pathway 2: Immuno-modulators (page 4)

Consider biologics if severe active Crohn's disease (HBI $\geq 8-9$ or CDAI ≥ 300)

CROHN'S DISEASE – Pathway 3: High Cost Drugs (page 5)

MAINTAINING REMISSION

Discuss options of treatment and no treatment, including the risk of inflammatory exacerbations and side effects

No treatment

- Agree follow-up plan and frequency
- Give advice on symptoms that may suggest a relapse and require medical attention e.g.
 - Unexplained weight loss
 - Abdominal pain
 - Diarrhoea
 - General ill-health
- Emphasise importance of not smoking

After surgery i.e. ileocolonic Crohn's disease with complete macroscopic resection and no residual disease:

- Consider **azathioprine with up to 3 months metronidazole**
- Consider **azathioprine alone** (if metronidazole not tolerated)

Do not offer: biologics or budesonide

If residual active disease (see 'Inducing remission' above)


Immuno-modulator treatment

Do not offer: conventional glucocorticosteroids or budesonide

CROHN'S DISEASE – Pathway 2: Immuno-modulators (page 4)

SWL IBD Pathway CROHN'S DISEASE - Pathway 2: Immuno-modulators


(References: 5, 8, 9, 10, 11, 12, 13 with local agreements) (09/09/2020)



Dosing table [ref 8,9,10,11,12]	TPMT normal activity (wild type)	TPMT intermediate activity (heterozygous)	TPMT low to no activity (homozygous)
Azathioprine (AZA)	100% dose [2.0 - 2.5 mg/kg] ^{8,9,11}	50% dose ^{8,9,11} [1.0 - 1.25 mg/kg]	Not recommended ¹²
Low dose azathioprine + allopurinol (100mg daily)	Initially 25% ^{8,12} of monotherapy dose [0.50 - 0.625mg/kg] Up to 50% ¹¹ of monotherapy dose [1.0 - 1.25mg/kg]	Initially 25% ^{8,12} of monotherapy dose [0.25 - 0.313mg/kg] Up to 50% ⁸ of monotherapy dose [0.5 - 0.625mg/kg]	Not recommended ¹²
6-mercaptopurine (half of AZA dose)	100% dose [1.0 - 1.5 mg/kg] ^{8,9,11}	50% dose ¹¹ [0.5 - 0.75mg/kg]	Not recommended ¹²
Low dose 6-mercaptopurine + allopurinol (100mg daily)	Initially 25% ^{8,12} of monotherapy dose [0.25 - 0.375mg/kg] Up to 50% ¹¹ of monotherapy dose [0.5 - 0.75mg/kg]	Initially 25% ^{8,12} of monotherapy dose [0.125 - 0.188mg/kg] Up to 50% ¹¹ of monotherapy dose [0.25 - 0.375mg/kg]	Not recommended ¹²

SWL IBD Pathway- CROHN'S DISEASE - Pathway 3: High Cost Drugs

(Reference: NICE¹⁴ with local agreements) (09/09/2020)



Pathway 3A: Severe active Crohn's Disease

Step 1: Use least expensive drug as a planned course of treatment (note 1,2)

1st choice: Adalimumab (TA187) (note 4) or
Infliximab biosimilar (TA187) (note 4) or
2nd choice: Ustekinumab (TA456)

If both TNF-alpha inhibitors contra-indicated or not tolerated (note 3)

1st choice: Ustekinumab (TA456) or
2nd choice: Vedolizumab (TA352)

At any point before 12 months of treatment has passed, has treatment failed (including the need for surgery)? (note 4,5,6)

Step 2: **1st choice:** Adalimumab (note 4) or
Infliximab biosimilar (note 4) or
2nd choice: Ustekinumab (TA456) or
3rd choice: Vedolizumab (TA352)

At any point before 12 months of treatment has passed, has treatment failed (including the need for surgery)? (note 4,5,6)

Step 3: **1st choice:** Adalimumab (note 4) or
Infliximab biosimilar (note 4) or
2nd choice: Ustekinumab or
3rd choice: Vedolizumab

At any point before 12 months of treatment has passed, has treatment failed (including the need for surgery)? (note 4,5,6)

Discontinue high cost drug treatment (note 8)

At 12 months after the start of treatment, reassess the disease (note 7). Is there evidence of ongoing active disease?

Yes Is the patient in stable clinical remission?

Discuss the risks and benefits of continued treatment. Is a trial withdrawal considered appropriate?

Trial withdrawal from drug used. Restart treatment if patient relapses after treatment is stopped

Maintain treatment & reassess patient at least every 12 months (note 5,6)
If appropriate move to **next step** in treatment pathway if response is lost at any point during therapy

Note 1: Choose **ONE** drug per step (note that step 2 is optional) before moving onto the next step due to primary or secondary treatment failure.

If there is more than one NICE approved treatment available, NICE recommends a discussion between the responsible clinician and the patient about the advantages and disadvantages of each treatment (considering therapeutic need and likely adherence to treatment). If more than one treatment option is suitable, the least expensive will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs.

The SWL drug choices in this algorithm are based on cost (using list price or nationally (NICE) / locally (LPP) agreed contract prices).

Note 4: If treatment failure with adalimumab or infliximab, an alternative TNF-alpha inhibitor may be chosen from the same step, if considered clinically appropriate. This is restricted to **ONE** switch within the TNF-alpha inhibitor class only.

Note 5: Treatment response

At 12 weeks (adalimumab), 6 weeks (infliximab), 14 weeks (vedolizumab), 16 weeks (ustekinumab) after the start of treatment, people should have their disease reassessed to determine whether patients are responding adequately and if ongoing treatment is still appropriate. Treatment should only be continued if there is clear evidence of response to drug treatment defined by a decrease in Harvey-Bradshaw Index by ≥ 3 points or CDAI by ≥ 70 points.

Note 6: Temporary dose escalation is commissioned for patients with secondary treatment failure as follows:

- Adalimumab: 40mg / week for 3 months or 80mg / 2 weeks for 3 months
- Infliximab: 5mg/kg / 4-6 weeks for 3 doses or 10mg/kg / 8 weeks for 3 doses
- Ustekinumab: 90mg / 8 weeks for 4 months
- Vedolizumab: 300mg / 4 weeks for 3 months, ONLY if no alternative drug options exist. Dose escalation is not commissioned if alternative drug options can be used.

For details and subsequent dose escalation requests see page 14.

Clinicians and commissioners should refer to the relevant technology appraisal and SPC for each drug for further information about their eligibility and prescription.

Note 2: Consider alternative drug from the same step if treatment had to be stopped due to an adverse event in patients who:


- had responded to CCG approved drug **OR**
- treatment response was not yet assessed i.e. before 12 weeks (adalimumab), 6 weeks (infliximab), 14 weeks (vedolizumab), 16 weeks (ustekinumab).

Note 3: For contraindications, cautions and information on malignancies see page 13

SWL IBD Pathway CROHN'S DISEASE - Pathway 3: High Cost Drugs

(Reference: NICE ¹⁴ with local agreements) (09/09/2020)

Pathway 3B: Active fistulising Crohn's Disease



Advice: Discuss the condition, associated symptoms, treatment options and monitoring with the patient and/or their parent or carer
Give advice on nature, frequency and severity of side effects of treatment and colorectal cancer risk

INDUCING REMISSION

Mild to moderate ulcerative colitis

For patients with a mild to moderate first presentation or inflammatory exacerbation

Acute severe ulcerative colitis

Proctosigmoiditis or left-sided ulcerative colitis:

- Topical aminosalicylate

If remission not achieved within 4 weeks:

- add oral aminosalicylate (high dose) **or**
- switch to oral aminosalicylate (high dose) + topical corticosteroid (time limited)

If further treatment needed:

- Stop topical treatments and offer oral aminosalicylate + oral corticosteroid (time limited e.g. prednisolone 40mg OD for 8 weeks)

If patient declines topical treatments:

- High dose oral aminosalicylate alone (not as effective as topical aminosalicylate)
- Add time limited course of oral corticosteroid (if remission not achieved within 4 weeks)

If patient cannot tolerate aminosalicylates:

- Topical or oral corticosteroid (time limited course)

Extensive disease:

- Topical aminosalicylate + high-dose oral aminosalicylate

If remission not achieved within 4 weeks:

- Switch to high dose oral aminosalicylate + oral corticosteroid (time limited)

If patient cannot tolerate aminosalicylates:

- Oral corticosteroid (time limited course)

Proctitis:

- Topical aminosalicylate

If remission not achieved within 4 weeks:

- Add oral aminosalicylate

If further treatment needed:

- Add topical or oral corticosteroid (time limited course e.g. prednisolone 40mg OD for 8 weeks)

If patient declines topical aminosalicylate:

- Oral aminosalicylate (not as effective as topical aminosalicylate)

If remission not achieved within 4 weeks:

- Add topical or oral corticosteroid (time limited course e.g. prednisolone 40mg OD for 8 weeks)

If patient cannot tolerate aminosalicylates:

- Topical or oral corticosteroid (time limited course e.g. prednisolone 40mg OD for 8 weeks)

Admit to hospital for treatment by gastroenterologist and colorectal surgeon (and paediatrician or obstetrics & gynaecology, paediatrician team if necessary)

Assess likelihood of needing surgery

ULCERATIVE COLITIS - Pathway 3: High Cost Drugs (page 10)

MAINTAINING REMISSION

Mild to moderate ulcerative colitis

Acute severe ulcerative colitis (single episode)

Proctitis or proctosigmoiditis:

- Topical aminosalicylate (daily or intermittent), with or without an oral aminosalicylate **or**
- Oral aminosalicylate alone (less effective)

Left-sided or extensive ulcerative colitis:

- Oral aminosalicylate (give a low maintenance dose in adults)

For patients with ≥ 2 inflammatory exacerbations in a 12 months period requiring systemic corticosteroids or in whom remission is not maintained with aminosalicylates

Consider immuno-modulators

For patients who decline, cannot tolerate or have a contraindication to immuno-modulators:

- Consider oral aminosalicylate


ULCERATIVE COLITIS - Pathway 2: Immuno-modulators (page 9)

Monitoring:

Adults: Offer colonoscopic surveillance according to NICE CG118⁶. Assess the risk of fragility fractures according to NICE CG146⁷.

SWL IBD Pathway ULCERATIVE COLITIS - Pathway 2: Immuno-modulators


(References: 8, 9, 10, 11, 12, 15, 16 with local agreements) (09/09/2020)



Dosing table [ref 8, 9, 10, 11, 12]	TPMT normal activity (wild type)	TPMT intermediate activity (heterozygous)	TPMT low to no activity (homozygous)
Azathioprine (AZA)	100% dose [2.0 - 2.5 mg/kg] ^{8,9,11}	50% dose ^{8,9,11} [1.0 - 1.25 mg/kg]	Not recommended ¹²
Low dose azathioprine + allopurinol (100mg daily)	Initially 25% ^{8,12} of monotherapy dose [0.50 - 0.625mg/kg] Up to 50% ¹¹ of monotherapy dose [1.0 - 1.25mg/kg]	Initially 25% ^{8,12} of monotherapy dose [0.25 - 0.313mg/kg] Up to 50% ⁸ of monotherapy dose [0.5 - 0.625mg/kg]	Not recommended ¹²
6-mercaptopurine (half of AZA dose)	100% dose [1.0 - 1.5 mg/kg] ^{8,9,11}	50% dose ¹¹ [0.5 - 0.75mg/kg]	Not recommended ¹²
Low dose 6-mercaptopurine + allopurinol (100mg daily)	Initially 25% ^{8,12} of monotherapy dose [0.25 - 0.375mg/kg] Up to 50% ¹¹ of monotherapy dose [0.5 - 0.75mg/kg]	Initially 25% ^{8,12} of monotherapy dose [0.125 - 0.188mg/kg] Up to 50% ¹¹ of monotherapy dose [0.25 - 0.375mg/kg]	Not recommended ¹²


SWL IBD Pathway ULCERATIVE COLITIS - Pathway 3: High Cost Drugs

(Reference: NICE¹⁷ with local agreements) (09/09/2020)



No

Pathway 3A: Moderately to severely active ulcerative colitis managed in outpatients



Pathway 3B: Acute exacerbation of severely active ulcerative colitis with hospitalisation

Acute exacerbation of severely active ulcerative colitis
 Severely active ulcerative colitis requires hospitalisation and urgent consideration for surgery and is defined by the following symptoms (Truelove & Witts)

- Bowel movements ≥ 6 plus at least one of the features of systemic upset (marked with *)
- Blood in stools – visible blood
- *Pyrexia (temperature greater than 37.8C)
- *Pulse > 90 bpm
- *Anaemia
- *ESR > 30 mm/hr

This clinical definition corresponds to

- Mayo score ≥ 9
- Partial Mayo score ≥ 6


Note 8: Funding requests for treatment outside this commissioned pathway can be made via the Individual Funding Request (IFR) process to the relevant commissioning organisation (see www.swlmcg.nhs.uk for IFR policy and application form)

Note 9: Contraindications to ciclosporin IV:

- Hypersensitivity to the active substance or to any of the excipients listed in the Summary of Product Characteristics (SPC)
- Combination with products containing *hypericum perforatum*
- Combination with medicines that are substrates for the multidrug efflux transporter P-glycoprotein or the organic anion transporter proteins (OATP) and for which elevated plasma concentrations are associated with serious and/or life-threatening events, e.g. bosentan, dabigatran etexilate and aliskiren

NOTE: This information is not exhaustive. Please also consult the SPC for the respective drug prior to prescribing for up to date prescribing information, including detailed information on adverse effects, drug interactions, cautions and contraindications (available via www.medicines.org.uk).

Has the patient received 72 hours of IV corticosteroids and urgent surgical intervention is being considered because of worsening symptoms
AND
 Is the patient contraindicated to ciclosporin or is ciclosporin considered to be clinically inappropriate? (note 8 & 9)



SWL IBD Pathway High Cost Drug

Contraindications, cautions and information about malignancies (09/09/2020)

NOTE: The information in tables 1 and 2 is not exhaustive. Please also consult the Summary of Product Characteristics (SPC) for the respective drugs prior to prescribing for up to date prescribing information, including detailed information on adverse effects, drug interactions, cautions and contraindications (available via www.medicines.org.uk).

Table 1: Contraindications

TNF-alpha inhibitors ^{18,19,20}	Ustekinumab ²¹	Vedolizumab ²²	Tofacitinib ²³
<ul style="list-style-type: none"> Hypersensitivity to active substance or excipients Active TB and other severe infections (sepsis, abscesses) and opportunistic infections Moderate to severe heart failure (NYHA class III/IV) 	<ul style="list-style-type: none"> Hypersensitivity to active substance or excipients Clinically important, active infection (eg active TB) 	<ul style="list-style-type: none"> Hypersensitivity to active substance or excipients Active severe infections such as TB, sepsis, CMV, listeriosis, and opportunistic infections such as PML 	<ul style="list-style-type: none"> Hypersensitivity to active substance or excipients Active TB, serious infections (e.g. sepsis) or opportunistic infections Severe hepatic impairment Pregnancy and lactation

Table 2: Cautions

TNF-alpha inhibitors ^{18,19,20}	Ustekinumab ²¹	Vedolizumab ²²	Tofacitinib ²³
<ul style="list-style-type: none"> Infections (impaired lung function) Hepatitis B reactivation Demyelinating diseases Malignancies – lymphomas Non-melanoma skin cancer (history of prolonged immuno-modulator therapy, PUVA) Mild heart failure (NYHA class I/II) Autoimmune processes (Lupus) 	<ul style="list-style-type: none"> Chronic infections or history of recurrent infections Malignancies – no studies Non-melanoma skin cancer (> 60 years of age, history of prolonged immuno-modulator therapy, PUVA) 	<ul style="list-style-type: none"> No identified systemic immunosuppressive activity but effects on systemic immune system function in patients with IBD not known The risk of malignancy is increased in patients with ulcerative colitis and Crohn's disease. Immuno-modulatory medicinal products may increase the risk of malignancy Malignancies not listed as side-effect 	<ul style="list-style-type: none"> Serious infections TB Viral reactivation Malignancy and lymphoproliferative disorder Non-melanoma skin cancer Interstitial lung disease GI perforations Liver enzymes Hypersensitivity Laboratory parameters (lymphocytes, neutrophils, haemoglobin, lipids)- see SPC for details

Table 3: immuno-modulator therapies to use or avoid in IBD patients with a history of cancer ²⁴

Type of cancer	Avoid	Use with caution	Can be used
Lymphoma	Thiopurines	Anti-TNF, methotrexate, steroids	
Acute myeloid leukaemia and severe myelodysplastic disorders	Thiopurines	Anti-TNF	Methotrexate, steroids
Melanoma	Anti-TNF	Thiopurines, steroids	Methotrexate
Non-melanoma skin cancer	Thiopurines	Anti-TNF, steroids	Methotrexate
Urinary tract cancer	Thiopurines	Anti-TNF	Methotrexate, steroids
Other tumours		Thiopurines, anti-TNF	Methotrexate, steroids

Table 4: Risk of cancer occurrence ²⁴

Risk	Organ/type of cancer
Low [< 10%]	Incidental asymptomatic renal tumour Lymphomas Testicle
Intermediate [11-25%]	Uterine body Colon Prostate Breast
High [>25%]	Bladder Sarcoma Melanoma and non-melanoma skin cancer Myeloma Symptomatic renal carcinoma

Dose escalation strategy

(References: 18, 19, 20, 21, 22, 23 with local agreements) (09/09/2020)

SWL dose escalation agreement for patients with secondary treatment failure:

- (1) 1st course of temporary dose escalation as per table below [Blueteq application[#]]
- (2) De-escalation to standard dose after 1st temporary escalation course
- (3) 2nd course of temporary dose escalation if relapse occurs after > 1 month as per table below [Blueteq application[#]]
- (4) De-escalation to standard dose after 2nd temporary escalation course
- (5) Continuous (up to 1 year) dose escalation if rapid relapse occurs (< 1 month) or if relapse occurs after 2nd temporary course [Blueteq application[#]] *
- (6) After 1 year de-escalation to standard dose (if in remission) or provide evidence for active residual disease and agreement following multi-Trust MDT discussion (requires submission of notes of multi-Trust MDT discussion and agreement. Participants must include gastroenterologists, colorectal surgeon, clinical nurse specialist, dietitian, pharmacist, pathologist and radiologist with special interest in gastroenterology as per IBD standards, standard A1 and A2²⁵)**

* Not applicable to vedolizumab

Golimumab dose escalation from 50mg to 100mg every 4 weeks in patients under 80kg is routinely commissioned and does not require a dose escalation Blueteq application

Drug	Crohn's disease	Ulcerative colitis
Adalimumab ¹⁸	40mg every week for 3 months or 80mg every 2 weeks for 3 months	40mg every week for 3 months or 80mg every 2 weeks for 3 months
Infliximab ¹⁹	5mg/kg every 6 weeks for 3 doses or 5mg/kg every 4 weeks for 3 doses or 10mg/kg every 8 weeks for 3 doses	If low drug concentrations (<5micrograms/ml) or antibodies to infliximab: 5mg/kg every 6 weeks for 3 doses or 5mg/kg every 4 weeks for 3 doses or 10mg/kg every 8 weeks for 3 doses (unlicensed; SWL local agreement)
Golimumab ²⁰		100mg every 4 weeks in patients <80kg [#] (routinely commissioned)
Ustekinumab ²¹	90mg every 8 weeks for 4 months	90mg every 8 weeks for 4 months
Vedolizumab ²²	300mg every 4 weeks for 3 months if no alternative drug options are considered appropriate by the clinician. Dose escalation is not commissioned if alternative drug options can be used.	300mg every 4 weeks for 3 months if no alternative drug options are considered appropriate by the clinician. Dose escalation is not commissioned if alternative drug options can be used.
Tofacitinib ²³		10mg twice daily for 4 months

SWL IBD Pathway - References

(09/09/2020)

References:

1. NICE DG 11 – Faecal calprotectin diagnostic tests for inflammatory bowel disease (Oct 2013)
2. Inflammatory Bowel Disease Toolkit. RCGP (see: <https://www.rcgp.org.uk/clinical-and-research/resources/toolkits/inflammatory-bowel-disease-toolkit.aspx>)
3. SEL APC. Primary & Secondary Care Inflammatory Bowel Disease Pathway (Feb 2018).
4. Christian M. ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 1: Initial diagnosis, monitoring of known IBD, detection of complications. *Journal of Crohn's and Colitis* 2019; 13(2): 144–164.
5. NICE Guideline 129 (NG129), Crohn's disease: management. May 2019.
6. NICE Clinical Guideline 118 (CG118). Colorectal cancer prevention: colonoscopic surveillance in adults with ulcerative colitis, Crohn's disease or adenomas. March 2011.
7. NICE Clinical Guidelines 146 (CG146). Osteoporosis: assessing the risk of fragility fracture. August 2012, last updated February 2017.
8. Ansari *et al.* Influence of xanthine oxidase on thiopurine metabolism in Crohn's disease. *Aliment Pharmacol Ther* 2008; 28: 749-757.
9. Gomollon *et al.* 3rd European Evidence-based Consensus on the Diagnosis and Medical Management of Crohn's Disease 2016: Part 1: Diagnosis and Management. *Journal of Crohn's and Colitis* 2017: 3-25.
10. Harbord *et al.* Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 2: Current Management. *Journal of Crohn's and Colitis* 2017, 769-784.
11. Warner *et al.* A practical guide to thiopurine prescribing and monitoring in IBD. *Frontline Gastroenterology* 2016; 0: 1-6.
12. Inflammatory Bowel Disease Handbook. 1st edition. Guy's and St. Thomas's IBD centre, 2014.
13. Gionchette P *et al.* 3rd European Evidence-based Consensus on the Diagnosis and Medical Management of Crohn's Disease 2016: Part 2: Surgical Management and Special Situations. *Journal of Crohn's and Colitis* 2017: 135-149.
14. NICE Crohn's disease pathway (see: <https://pathways.nice.org.uk/pathways/crohns-disease>)
15. NICE Guideline 130 (NG130), Ulcerative colitis: management. May 2019.
16. Travis S P L *et al.* Predicting outcome in severe ulcerative colitis. *Gut* 1996; 38(6):905–910.
17. NICE Ulcerative Colitis pathway (see: <https://pathways.nice.org.uk/pathways/ulcerative-colitis>)
18. www.medicines.org.uk. Accessed 01/06/2019; Humira® (adalimumab) Summary of Product Characteristics (SPC) last updated 29/05/2019.
19. www.medicines.org.uk. Accessed 01/06/2019; Remicade® (infliximab) Summary of Product Characteristics (SPC) last updated 01/04/2019.
20. www.medicines.org.uk. Accessed 01/06/2019; Simponi® (golimumab) Summary of Product Characteristics (SPC) last updated 28/03/2019.
21. www.medicines.org.uk. Accessed 01/06/2019; Stelara® (ustekinumab) Summary of Product Characteristics (SPC) last updated 16/01/2019.
22. www.medicines.org.uk. Accessed 01/06/2019; Entyvio® (vedolizumab) Summary of Product Characteristics (SPC) last updated 07/03/2019.
23. www.medicines.org.uk. Accessed 01/06/2019; Xeljanz® (tofacitinib) Summary of Product Characteristics (SPC) last updated 16/11/2018.
24. Annese V *et al.* ECCO Guideline/Consensus Paper, European Evidence-based Consensus: Inflammatory Bowel Disease and Malignancies. *Journal of Crohn's and Colitis* 2015: 945-965.
25. IBD Standards, 2013 update. Standard for the healthcare of people who have inflammatory bowel disease (IBD) (see: <http://s3-eu-west-1.amazonaws.com/files.crohnsandcolitis.org.uk/Publications/PPR/ibd-standards.pdf>)

SWL IBD Pathway – Version control

(09/09/2020)

Version number	Amendments made	Date of approval
0		26 May 2011
1.0	<p>Include approved recommendations from South West London Biologics Care Pathway Review for Inflammatory Bowel Disease (IBD) (8 Feb 2017):</p> <ul style="list-style-type: none"> • Local agreement (Crohn's disease) – allow use of alternative TNF-alpha inhibitor (adalimumab or infliximab) if appropriate (step 2) • Local agreement (ulcerative colitis) – allow use of alternative TNF-alpha inhibitor (adalimumab, infliximab or golimumab (if high BMI and >100kg)) if appropriate (step 2) 	27 Feb 2017
2.0	<p>Include approved recommendations from SWL IBD network meeting (28 Jun 2017):</p> <ul style="list-style-type: none"> • Preferred drug choices • Add existing agreements on dose escalation • Add contraindications and information on cancer risk • Improved pathway presentation • Crohn's disease: <ul style="list-style-type: none"> ➢ Ustekinumab (NICE TA 456) in step 1 (only if anti-TNF contraindicated/not tolerated (local agreement)) and in step 3 ➢ New local agreement: vedolizumab step 4 ➢ Add dose escalation with ustekinumab in line with license and NICE TA 	01 Nov 2017
2.1	<p>Amend note 1 to clarify that a discussion between the responsible clinician and the patient should take place about the advantages and disadvantages of each treatment (considering therapeutic need and likely adherence to treatment) and if more than one treatment option is suitable, the least expensive will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs.</p>	11 Jan 2018
3.0	<p>Include approved recommendations from SWL IBD network meeting (12 Jul 2018):</p> <ul style="list-style-type: none"> • Change presentation to clarify that step 2 is an optional step and not mandated (local agreement) • Include note 7- reference to IFR process • Crohn's disease: <ul style="list-style-type: none"> ➢ Change pathway presentation to clarify that TNF-alpha inhibitors are not currently commissioned after ustekinumab or vedolizumab • Ulcerative colitis: <ul style="list-style-type: none"> ➢ Change pathway presentation to clarify that TNF-alpha inhibitors are not currently commissioned after use of vedolizumab 	08 Oct 2018
4.0	<p>Include approved recommendations from SWL IBD network meeting (28 Feb 2019):</p> <ul style="list-style-type: none"> • Integration of pathways into one SWL IBD pathway including: <ul style="list-style-type: none"> ➢ Presenting with symptoms ➢ Inducing and maintaining remission ➢ Name change from "Drug pathway" to "High cost drug pathway" • Crohn's disease: <ul style="list-style-type: none"> ➢ Remove the following statement for vedolizumab: "if unable to use SC alternative" ➢ Include note 4 instead of "optional step 2" ➢ Add adalimumab and infliximab biosimilar as 1st choice options in all steps ➢ In the final step, add ustekinumab as an option • Ulcerative colitis: <ul style="list-style-type: none"> ➢ Include tofacitinib in step 1, 2 and 3 ➢ Include all treatment option in step 1,2 and 3 ➢ Include note 4 instead of "optional step 2" ➢ New dose escalation policy 	04 Oct 2019
4.1	<p>Include recommendations from St George's NHS Foundation Trust- Infliximab dose escalation (unlicensed) for Ulcerative Colitis (Dec 2019) following approval through SWL Trust Governance processes:</p> <p>Include infliximab dose escalation (5mg/kg every 6 weeks for 3 doses or 5mg/kg every 4 weeks for 3 doses or 10mg/kg every 8 weeks for 3 doses (unlicensed; SWL local agreement) in pathway 3A, note 6 and page 14 (dose escalation strategy) if low drug concentrations (<5 micrograms/ml) or antibodies to infliximab</p>	26 Mar 2020

SWL IBD Pathway – Version Control

(09/09/2020)

4.2	Add Ustekinumab (NICE TA633) to Ulcerative Colitis pathway: • Insert ustekinumab- 3 rd choice option to step 1, 2 and 3 • Move vedolizumab from 3 rd to 4 th choice option. • Include ustekinumab dose escalation (90mg / 8 weeks for 4 months)	9 Sept 2020
-----	---	-------------

Date of next review: September 2022 (or earlier if indicated)

ADDENDUM 1

(Approved by SWL Integrated Medicines Optimisation Committee on 20 July 2022)

NICE published TA792 (June 2022) - Filgotinib for treating moderately to severely active ulcerative colitis. This addendum aims to inform clinicians that filgotinib is available as a treatment option in line with NICE recommendations and local agreements. Based on its relative cost* filgotinib is commissioned as follows:

Pathway 3A

- Step 1 – 1st choice option
- Step 2 – 1st choice option
- Step 3 – 1st choice option

* The SWL drug choices in the “SWL Inflammatory Bowel Disease Pathway – Ulcerative Colitis” are based on cost (using list price or nationally (NICE) / locally (LPP) agreed contract prices).

ADDENDUM 2

(Approved by SWL Integrated Medicines Optimisation Committee on 21st December 2022)

NICE published TA828 (Oct 2022) – Ozanimod for treating moderately to severely active ulcerative colitis. This addendum aims to inform clinicians that ozanimod is available as a treatment option in line with NICE recommendations and local agreements. Based on its relative cost* ozanimod is commissioned as follows:

Pathway 3A

- Step 1 – 2nd choice option (only if infliximab is not suitable)
- Step 2 – 2nd choice option (infliximab is not suitable or already tried)
- Step 3 – 2nd choice option (infliximab is not suitable or already tried)